HVLab Vibrotactile Perception Meter:

Lewis, C.H. (1992) The HVLab Tactile Vibrometer, Proceedings from the United Kingdom Meeting of Human Responses to Vibration, ISVR, 28-30 September, 83-96.

Lindsell, C.J., Griffin, M.J. (1995). Signs and symptoms of disorders associated with hand-transmitted vibration in a group of dockyard workers. I.S.V.R. Contract Report No. 95/28, Institute of Sound and Vibration Research, University of Southampton.

Lindsell CJ (1997) Vibrotactile thresholds: Effect of contact forces and skin indentation, Proceedings of UK Group Meeting on Human Response to Vibration, Human Factors Research Unit, Southampton, UK, 17-19 September, 433-443.

Lindsell CJ and Griffin MJ (1998) Standardised diagnosis methods for assessing components of the hand-arm vibration syndrome. HSE Books, CRR197/98, ISBN 0717616401

Lindsell CJ and Griffin MJ (1999) Thermal thresholds, vibrotactile thresholds and finger systolic blood pressures in dockyard workers exposed to hand-transmitted vibration. International Archive of Occupational and Environmental Health, 72:377-386

Lindsell CJ and Griffin MJ (2002) Normative data for vascular and neurological tests of the hand-arm vibration syndrome. International Archive of Occupational and Environmental Health, 75:43-54

Morioka M and Griffin MJ (2005) Thresholds for the perception of hand-transmitted vibration: Dependence on contact area and contact location, Somatosensory & Motor Research 22(4):281-297

Whitehouse DJ, Morioka M and Griffin MJ (2006) Effect of contact location on vibrotactile thresholds at the fingertip, Somatosensory & Motor Research 23(1/2):73-81

Gu C and Griffin MJ (2007) Effect of frequency and contact location on vibrotactile perception thresholds at the foot. 11th International Conference on Hand-Arm Vibration,3-7 June 2007, Bologna, Italy.

Gu C (2007) The effect of surround contact force on vibrotactile perception thresholds at the big toe. The 42nd United Kingdom Conference on Human Responses to Vibration, held at ISVR, University of Southampton, Southampton, England, 10 - 12 September 2007

Morioka M, Whitehouse DJ and Griffin MJ (2008) Vibrotactile thresholds at the fingertip, volar forearm, large toe, and heel, Somatosensory & Motor Research 25(2):101-112

Seah SA and Griffin MJ (2008) Normal values for thermotactile and vibrotactile thresholds in males and females. Int Arch Occup Environ Health 81:535-543

Gu C and Griffin MJ (2009) Effect of contact area on thresholds for the perception of 160-Hz vibration at the big toe. The 44th United Kingdom Conference on Human Responses to Vibration, held at Loughborough University, UK, 7 - 9 September

Maeda S, Yonekawa Y, Kanada K and Takahashi Y (1998) Effect of push forces on vibrotactile thresholds measurement, 8th International Conference on Hand-Arm Vibration, 9-12 June, Umeå, Sweden.

McGeoch KL and Gilmour WH (2000) Cross sectional study of a workforce exposed to hand-arm vibration: with objective tests and the Stockholm workshop scales. British Medical Journal, 57:35-42

Wild P, Massin N, Lasfargues G, Baudin V, Unlu D, Donati P (2001) Vibrotactile perception thresholds in four non-exposed populations of working age. Ergonomics, 44(6):649-657

Lawson IJ and McGeoch KL (2003) A medical assessment process for a large volume of medicao-legal compensation claims for hand-arm vibration syndrome. Occupational Medicine. 53:302-308

Daud R, Maeda S, Kameel NNM, Ripin MY, Bakrun N, Zein RM, Kido M, and Higuchi K (2004) A pilot study of reference vibrotactile perception threshold on the fingertip obtained with Malaysian healthy people using ISO 13091-1 equipment. Industrial Health, 42:189-195

Poole K and Mason H (2007) Relationship between self-reported upper limb disability and quantitative tests in hand-arm vibration syndrome. Disability and Rehabilitation, 29(5):359-366

Lundstrom R, Nilsson T, Hagberg M and Burstrom L (2008) Grading of sensorineural disturbances according to a modified Stockholm workshop scale using self-reports and QST. International Archive of Occupational and Environmental Health, 81:553-557

Hohne A, Stark C, Bruggemann GP (2009) Plantar pressure distribution in gait is not affected by targeted reduced plantar cutaneous sensation. Clinical Biomechanics, 24:308-313